

IN THE CLAIMS:

Claims 1-32 (cancelled)

33. (previously presented) A method of preparing a composition for the treatment of dermatological disorders, comprising the steps of: filtering coal tar to remove impurities by filter compression and preparing a composition wherein the composition includes the filtered coal tar.

34. (previously presented) A method as in Claim 33, further comprising the step of adding zinc pyrithione into the composition.

35. (previously presented) A method as in Claim 33, wherein the coal tar is filtered by being fed through a compressed charcoal filter.

36. (previously presented) A method as in Claim 35, wherein the coal tar is fed through the filter at approximately 17 to 19 psi.

37. (previously presented) A method as in Claim 35, wherein the coal tar is fed through the filter at approximately 18 psi.

38. (previously presented) A method as in Claim 35, further comprising the step of using compressed air to force the coal tar through the compressed charcoal filter.

39. (previously presented) A method as in Claim 35, further comprising the step of leaving the coal tar in the compressed charcoal filter for 8 hours.

40. (previously presented) A method as in Claim 35, further comprising the step of boiling a fluid that has been passed through the compressed charcoal filter.

41. (previously presented) A method as in Claim 40, wherein the fluid that has been passed through the compressed charcoal filter is boiled for 5 minutes.

42. (previously presented) A method as in Claim 40, wherein the boiled, filtered fluid is allowed to cool to room temperature.

43. (previously presented) A method as in Claim 40, further comprising the step of refiltering a top layer of the boiled, filtered fluid.

44. (previously presented) A method as in Claim 43, wherein refiltering is through a nylon mesh.

45. (previously presented) A method as in Claim 33, where a surfactant is added to the formulation.

46. (previously presented) A method as in Claim 45, wherein the surfactant is an ionic surfactant.

47. (previously presented) A method as in Claims 45, wherein the surfactant is sodium lauryl sulphate.

48. (previously presented) A method as in any Claim 33, wherein a carrier is added to the formulation.

49. (previously presented) A method as in Claim 48, wherein the carrier is isopropyl myristate.

50. (previously presented) A method as in Claim 48, wherein the carrier is ethyl alcohol.

51. (previously presented) A method as in Claim 33, further comprising the step of placing the composition in a container selected from the group consisting of a spray container and an aerosol container.

52-65. (Cancelled)

Please add the following new claim:

66. (new) A method as in Claim 35, wherein a plurality of carriers are added to the formulation.